

# Nitro-functionalized Bis(pyrazolate) Metal-Organic Frameworks as Carbon Dioxide Capture Materials under Ambient Conditions

Mosca N., Vismara R., Fernandes J., Tuci G., Di Nicola C., Domasevitch K., Giacobbe C., Giambastiani G., Pettinari C., Aragones-Anglada M., Moghadam P., Fairen-Jimenez D., Rossin A., Galli S.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

---

## Abstract

© 2018 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim The metal-organic frameworks (MOFs)  $M(\text{BPZNO}_2)$  ( $M=\text{Co}, \text{Cu}, \text{Zn}$ ;  $\text{H}_2\text{BPZNO}_2=3\text{-nitro-4,4'-bipyrazole}$ ) were prepared through solvothermal routes and were fully investigated in the solid state. They showed good thermal stability both under a  $\text{N}_2$  atmosphere and in air, with decomposition temperatures peaking up to 663 K for  $\text{Zn}(\text{BPZNO}_2)$ . Their crystal structure is characterized by 3D networks with square ( $M=\text{Co}, \text{Zn}$ ) or rhombic ( $M=\text{Cu}$ ) channels decorated by polar  $\text{NO}_2$  groups. As revealed by  $\text{N}_2$  adsorption at 77 K, they are micro-mesoporous materials with BET specific surface areas ranging from 400 to 900  $\text{m}^2 \text{g}^{-1}$ . Remarkably, under the mild conditions of 298 K and 1.2 bar,  $\text{Zn}(\text{BPZNO}_2)$  adsorbs 21.8 wt %  $\text{CO}_2$  (4.95  $\text{mmol g}^{-1}$ ). It shows a Henry  $\text{CO}_2/\text{N}_2$  selectivity of 15 and an ideal adsorbed solution theory (IAST) selectivity of 12 at  $p=1$  bar. As a  $\text{CO}_2$  adsorbent, this compound is the best-performing MOF to date among those bearing a nitro group as a unique chemical tag. High-resolution powder X-ray diffraction at 298 K and different  $\text{CO}_2$  loadings revealed, for the first time in a  $\text{NO}_2$ -functionalized MOF, the insurgence of primary host-guest interactions involving the  $\text{C}(3)\text{-NO}_2$  moiety of the framework and the oxygen atoms of carbon dioxide, as confirmed by Grand Canonical Monte Carlo simulations. This interaction mode is markedly different from that observed in  $\text{NH}_2$ -functionalized MOFs, for which the carbon atom of  $\text{CO}_2$  is involved.

<http://dx.doi.org/10.1002/chem.201802240>

---

## Keywords

adsorption, host-guest interactions, metal-organic frameworks, N ligands, X-ray diffraction

## References

- [1] T. H. Oh, Renewable Sustainable Energy Rev. 2010, 14, 2697
- [2] Handbook of Porous Solids, Vol. 1-5 (Eds.: F. Schüth, K. S. W. Sing, J. Weitkamp), Wiley-VCH, Weinheim, 2002
- [3] H. Yang, Z. Xu, M. Fan, R. Gupta, R. B. Slimane, A. E. Bland, I. Wright, J. Environ. Sci. 2008, 20, 14
- [4] L. R. MacGillivray, C. M. Lukehart, Metal-Organic Framework Materials, Wiley, New York, 2014
- [5] B. Seyyedi, Metal-Organic Frameworks: A New Class of Crystalline Porous Materials, Lambert Academic, Saarbrücken, 2014

- [6] S. R. Batten, N. R. Champness, X.-M. Chen, J. Garcia-Martinez, S. Kitagawa, L. Öhrström, M. O'Keeffe, M. P. Suh, J. Reedijk, *Pure Appl. Chem.* 2013, 85, 1715–1724
- [7] P. Z. Moghadam, A. Li, S. B. Wiggins, A. Tao, A. G. P. Maloney, P. A. Wood, S. C. Ward, D. Fairen-Jimenez, *Chem. Mater.* 2017, 29, 2618–2625
- [8] M. P. Suh, H. J. Park, T. K. Prasad, D. W. Lim, *Chem. Rev.* 2012, 112, 782–835
- [9] A. H. Chughtai, N. Ahmad, H. A. Younus, A. Laypkovc, F. Verpoort, *Chem. Soc. Rev.* 2015, 44, 6804–6849
- [10] J. Liu, L. Chen, H. Cui, J. Zhang, L. Zhang, C.-Y. Su, *Chem. Soc. Rev.* 2014, 43, 6011–6061
- [11] A. Rossin, G. Tuci, L. Luconi, G. Giambastiani, *ACS Catal.* 2017, 7, 5035–5045
- [12] F.-Y. Yi, D. Chen, M.-K. Wu, L. Han, H.-L. Jiang, *ChemPlusChem* 2016, 81, 675–690
- [13] Z. Hu, B. J. Deibert, J. Li, *Chem. Soc. Rev.* 2014, 43, 5815–5840
- [14] P. Horcajada, C. Serre, G. Maurin, N. A. Ramsahye, F. Balas, M. Vallet-Regi, M. Sebban, F. Taulelle, G. Férey, J. Am. Chem. Soc. 2008, 130, 6774–6780
- [15] S. S. Han, D. Kim, D. H. Jung, S. Cho, S.-H. Choi, Y. J. Jung, *J. Phys. Chem. C* 2012, 116, 20254–20261
- [16] R. Banerjee, H. Furukawa, D. Britt, C. Knobler, M. O'Keeffe, O. M. Yaghi, *J. Am. Chem. Soc.* 2009, 131, 3875–3877
- [17] M. Zhang, Q. Wang, Z. Lu, H. Liu, W. Liu, J. Bai, *CrystEngComm* 2014, 16, 6287–6290
- [18] D. K. Maity, A. Halder, B. Bhattacharya, A. Das, D. Ghoshal, *Cryst. Growth Des.* 2016, 16, 1162–1167
- [19] Z. H. Rada, H. R. Abid, H. Sun, S. Wang, *J. Chem. Eng. Data* 2015, 60, 2152–2161
- [20] A. Khutia, C. Janiak, *Dalton Trans.* 2014, 43, 1338–1347
- [21] H. Reinsch, S. Waitschata, N. Stock, *Dalton Trans.* 2013, 42, 4840–4847
- [22] S. Biswas, T. Ahnfeldt, N. Stock, *Inorg. Chem.* 2011, 50, 9518–9526
- [23] J. Sim, H. Yim, N. Ko, S. B. Choi, Y. Oh, H. J. Park, S. Y. Park, J. Kim, *Dalton Trans.* 2014, 43, 18017–18024
- [24] J.-P. Zhang, Y.-B. Zhang, J.-B. Lin, X.-M. Chen, *Chem. Rev.* 2012, 112, 1001–1033
- [25] C. Pettinari, A. Tăbăcaru, S. Galli, *Coord. Chem. Rev.* 2016, 307, 1–31
- [26] V. Colombo, C. Montoro, A. Maspero, G. Palmisano, N. Masciocchi, S. Galli, E. Barea, J. A. R. Navarro, *J. Am. Chem. Soc.* 2012, 134, 12830–12843
- [27] V. Colombo, S. Galli, H. J. Choi, G. D. Han, A. Maspero, G. Palmisano, N. Masciocchi, J. R. Long, *Chem. Sci.* 2011, 2, 1311–1319
- [28] N. Masciocchi, S. Galli, V. Colombo, A. Maspero, G. Palmisano, B. Seyyedi, C. Lamberti, S. Bordiga, *J. Am. Chem. Soc.* 2010, 132, 7902–7904
- [29] S. Galli, N. Masciocchi, V. Colombo, A. Maspero, G. Palmisano, F. J. López-Garzón, M. Domingo-García, I. Fernández-Morales, E. Barea, J. A. R. Navarro, *Chem. Mater.* 2010, 22, 1664–1672
- [30] Z. R. Herm, B. M. Wiers, J. A. Mason, J. M. van Baten, M. R. Hudson, P. Zajdel, C. M. Brown, N. Masciocchi, R. Krishna, J. R. Long, *Science* 2013, 340, 960–964
- [31] F. Salles, G. Maurin, C. Serre, P. L. Llewellyn, C. Knofel, H. J. Choi, Y. Filinchuk, L. Oliviero, A. Vimont, J. R. Long, G. Férey, *J. Am. Chem. Soc.* 2010, 132, 13782–13788
- [32] J. P. Zhang, S. Kitagawa, *J. Am. Chem. Soc.* 2008, 130, 907–917
- [33] K. S. Park, Z. Ni, A. P. Côté, J. Y. Choi, R. Huang, F. J. Uribe-Romo, H. K. Chae, M. O'Keeffe, O. M. Yaghi, *Proc. Natl. Acad. Sci. USA* 2006, 103, 10186–10191
- [34] M. Eddaoudi, J. Kim, N. Rosi, D. Vodak, J. Wachter, M. O'Keeffe, O. M. Yaghi, *Science* 2002, 295, 469–472
- [35] C. Pettinari, A. Tăbăcaru, I. Boldog, K. V. Domasevitch, S. Galli, N. Masciocchi, *Inorg. Chem.* 2012, 51, 5235–5245
- [36] A. Tăbăcaru, C. Pettinari, I. Timokhin, F. Marchetti, F. Carrasco-Marín, F. J. Maldonado-Hódar, S. Galli, N. Masciocchi, *Cryst. Growth Des.* 2013, 13, 3087–3097
- [37] N. Mosca, R. Vismara, J. A. Fernandes, S. Casassa, K. V. Domasevich, E. Bailón-García, F. J. Maldonado-Hódar, C. Pettinari, S. Galli, *Cryst. Growth Des.* 2017, 17, 3854–3867
- [38] H. J. Choi, M. Dincă, J. R. Long, *J. Am. Chem. Soc.* 2008, 130, 7848–7850
- [39] V. A. Blatov, A. P. Shevchenko, D. M. Proserpio, *Cryst. Growth Des.* 2014, 14, 3576–3586
- [40] Empty volume was estimated with the software PLATON (A. L. Spek, *Acta Crystallogr. Sect. D* 2009, 65, 148–155) on an ordered model (built, respectively, in C2/c for Co-BPZNO·DMF and Zn-BPZNO·DMF and in P2/m for Cu-BPZNO·DMF) after removal of the clathrated solvent, supposing that the nitro group protrudes into the channels. In parenthesis, the empty volume calculated assuming that the nitro group is coplanar to the skeleton of the ligand
- [41] Pore volume was calculated from the ambient conditions empty volume, supposing that the nitro group protrudes into the channels. In parenthesis, the empty volume calculated assuming that the nitro group is coplanar to the skeleton of the ligand

- [42] F. Millange, C. Serre, G. Férey, *Chem. Commun.* 2002, 822–823
- [43] A. Demessence, J. R. Long, *Chem. Eur. J.* 2010, 16, 5902–5908
- [44] M. Dincă, A. F. Yu, J. R. Long, *J. Am. Chem. Soc.* 2006, 128, 8904–8913
- [45] M. Kandiah, M. H. Nilsen, S. Usseglio, S. Jakobsen, U. Olsbye, M. Tilset, C. Larabi, E. A. Quadrelli, F. Bonino, K. P. Lillerud, *Chem. Mater.* 2010, 22, 6632–6640
- [46] S. Galli, A. Maspero, C. Giacobbe, G. Palmisano, L. Nardo, A. Comotti, I. Bassanetti, P. Sozzani, N. Masciocchi, *J. Mater. Chem. A* 2014, 2, 12208–12221
- [47] E. J. Beckman, *Chem. Commun.* 2004, 1885–1888
- [48] L. Li, J. Yang, J. Li, Y. Chen, J. Li, *Microporous Mesoporous Mater.* 2014, 198, 236–246
- [49] L. Ding, A. O. Yazaydin, *Phys. Chem. Chem. Phys.* 2013, 15, 11856–11861
- [50] J. Yu, L.-H. Xie, J.-R. Li, Y. Ma, J. M. Seminario, P. B. Balbuena, *Chem. Rev.* 2017, 117, 9674–9754
- [51] B. Sreenivasulu, I. Sreedhar, P. Suresh, K. V. Raghavan, *Catal. Sci. Technol.* 2015, 49, 12641–12661
- [52] B. Bhattacharya, D. Ghoshal, *CrystEngComm* 2015, 17, 8388–8413
- [53] Y. Lu, M. Tonigold, B. Breidenkötter, D. Volkmer, J. Hitzbleck, G. Langstein, *Z. Anorg. Allg. Chem.* 2008, 634, 2411–2417
- [54] G. Férey, C. Serre, *Chem. Soc. Rev.* 2009, 38, 1380–1399
- [55] R. Vaidhyanathan, S. S. Iremonger, G. K. H. Shimizu, P. G. Boyd, S. Alavi, T. K. Woo, *Science* 2010, 330, 650–653
- [56] I. Boldog, A. N. Chernega, J. Sieler, K. V. Domasevitch, *Inorg. Chim. Acta* 2002, 338, 69–77
- [57] A. A. Coelho, *J. Appl. Crystallogr.* 2003, 36, 86–95
- [58] A. A. Coelho, *TOPAS-Academic V6*, 2016, <http://www.topas-academic.net>
- [59] Bond lengths and angles for the rigid body describing the ligand: C–C and C–N of the pyrazole ring 1.36 Å; exocyclic C–C 1.40 Å; C–H of the pyrazole ring 0.95 Å; C–NNO<sub>2</sub> 1.40 Å; N–O 1.23 Å; pyrazole ring internal and external bond angles 108 and 126°, respectively; angles at the nitrogen atom of the nitro group 120°. Bond lengths and angles for the rigid body describing the solvent: C–N 1.45 Å; C–H 0.95 Å; C=O 1.25 Å; angles at the nitrogen and amidic carbon atoms 120°; angles at the carbon atoms of the methyl groups 109.5°
- [60] A. A. Coelho, *J. Appl. Crystallogr.* 2000, 33, 899–908
- [61] Refined bond lengths in the final stages of Rietveld refinements: C–C 1.40–1.45 Å, C–C and C–N of the pyrazole ring 1.32–1.38 Å
- [62] See: [http://topas.dur.ac.uk/topaswiki/doku.php?id=background\\_straight\\_line\\_segments](http://topas.dur.ac.uk/topaswiki/doku.php?id=background_straight_line_segments)
- [63] R. W. Cheary, A. A. Coelho, *J. Appl. Crystallogr.* 1992, 25, 109–121
- [64] J. A. Mason, J. Oktawiec, M. K. Taylor, M. R. Hudson, J. Rodriguez, J. E. Bachman, M. I. Gonzalez, A. Cervellino, A. Guagliardi, C. M. Brown, P. L. Llewellyn, N. Masciocchi, J. R. Long, *Nature* 2015, 527, 357–361
- [65] When comparing the TGA and VT-PXRD results, the reader must be aware of the fact that the thermocouple of the VT-PXRD setup is not in direct contact with the sample. Consequently, a slight difference in the temperature at which the same event is detected by the two techniques is unavoidable. The TGA temperatures are more reliable
- [66] X. Zhu, S. M. Mahurin, S.-H. An, C.-L. Do-Thanh, C. Tian, Y. Li, L. W. Gill, E. W. Hagaman, Z. Bian, J.-H. Zhou, J. Hu, H. Liu, S. Dai, *Chem. Commun.* 2014, 50, 7933–7936
- [67] X. Zhu, C. Tian, G. M. Veith, C. W. Abney, J. Dehaut, S. Dai, *J. Am. Chem. Soc.* 2016, 138, 11497–11500
- [68] J. Schell, N. Casas, R. Pini, M. Mazzotti, *Adsorption* 2012, 18, 49–65
- [69] M. Brunelli, A. N. Fitch, *J. Synchrotron Radiat.* 2003, 10, 337–339
- [70] To build up the rigid body modelling CO<sub>2</sub>, the following bond distances and angles were adopted: C=O, 1.21 Å; O=C=O, 180°
- [71] A. K. Rappe, C. J. Casewit, K. S. Colwell, W. A. Goddard III, W. M. Skiff, *J. Am. Chem. Soc.* 1992, 114, 10024–10035
- [72] J. J. Potoff, J. I. Siepmann, *AIChE J.* 2001, 47, 1676–1682
- [73] *Materials Studio 5.0*, Accelrys Software Inc., San Diego, 2009
- [74] D. Fairen-Jimenez, Y. J. Colón, O. K. Farha, Y.-S. Bae, J. T. Hupp, R. Q. Snurr, *Chem. Commun.* 2012, 48, 10496–10498
- [75] W. Bury, D. Fairen-Jimenez, M. B. Lalonde, R. Q. Snurr, O. K. Farha, J. T. Hupp, *Chem. Mater.* 2013, 25, 739–744